

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING**

# TARGET

**DUST TREATMENT FOR INSECT CONTROL**

Active constituent:  
10g/kg Permethrin 25;75

A ready to use dust treatment for the control of cockroaches, ants, fleas, silverfish, bedbugs, mushroom flies, potato moths, hide beetles and European Wasps as per the directions for use.

**PTC  
Prescription  
Certified  
Treatment**

---

**PCT** INTERNATIONAL PTY LTD

**Pest Control Technologies International Pty Ltd  
PO Box 6078 Acacia Ridge Delivery Centre, Acacia Ridge Q. 4110**

**Freecall Number 1800 630 877**

™Registered Trademark of PCT International, Brisbane, Queensland, Australia

# TARGET

## DUST TREATMENT FOR INSECT CONTROL DIRECTIONS FOR USE

Situation		Target Insects	State	Rate	Critical Comments
Domestic, institutional and industrial premises	Food preparation and storage areas	Cockroaches	All States	10g per m2 applied with a powder blower	Blow powder into all cracks and crevices. Pay particular attention to warm, dark corners, around cookers, fireplaces, radiators, stoves, pipes, rubbish chutes and places where liquid sprays may not be used.
		Silverfish			Sprinkle the powder along runs, on and around nests. If ants are invading domestic premises apply powder to the routes by which they enter. Repeat as necessary.
	Ants	Dust powder over floor, brushing into carpet pile and blowing into areas where flea larvae develop.			
	Fleas	Apply powder to bedsteads, bedsprings, mattresses, floor coverings, unholstered furniture, cracks in walls, behind torn wallpaper, joints in woodwork and other harborages.			
	Living and sleeping areas	Bedbugs			
Skin and Hide preparations		Hide Beetles (Dermestesator)			a) Place skins on drying racks and apply dust using a manual or motorised duster. b) Dust exposed edges of dried, bailed skins.
Mushroom Beds		Mushroom Sciand flies			Dust mushroom beds no more frequently than twice a week to control adult and larvae flies. Discontinue dusting when mushrooms reach the size of a 10 cent piece.
Seed Potatoes		Potato Moth (Phthorimaea operculella)		600g/tonne or 30g/50kg bag of seed potato	Dust thoroughly immediately after harvest while filling storage containers. Treated seed potatoes are to be used only for sowing, do not use for food or feed purposes. Treated seed potatoes should be stored apart from other potatoes and bags or containers should be clearly marked to identify that the contents have been treated.
Large Wasp Nests		European Wasps	NSW VIC WA TAS NT	125g to 250g per nest	Use hand duster and supply powder through entrance holes (125-250g/nest depending on size of nest and number of entrance holes).
					Treat at night whenever possible. Operator should wear clothing protective against the wasp including long-sleeved overalls buttoned at wrist and throat, gloves, hat, and bee-veil.

**NOT TO BE USED FOR ANY OTHER PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**  
WITHHOLDING PERIOD - FOR MUSHROOMS DO NOT APPLY LATER THAN 3 DAYS BEFORE HARVEST.

**PCT** INTERNATIONAL PTY LTD

**N.R.A. APPROVAL NO. 46589/1**