

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



**Talon**<sup>®</sup>  
Rodenticide pellets

syngenta.

**ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUM**

Ready to use bait for use in and around industrial, commercial, public services, agricultural and domestic buildings.

Controls rodents resistant to other anticoagulants, including warfarin, coumatetralyl and bromadiolone

**ONE FEED KILLS RATS AND MICE#**



**Syngenta Australia Pty Ltd**  
Level 1, 2-4 Lyonpark Road, Macquarie Park NSW 2113

**In a transport emergency dial 000, Police or Fire Brigade**  
**For specialist advice in an emergency only, call 1800 033 111 (24 hours)**

**APVMA Approval No: 33896/55730**

TM

## **DIRECTIONS FOR USE**

### **Restraints**

Place the ready-to-use bait in tamper-resistant bait stations

DO NOT remove pellets from bait station

DO NOT place bait stations in the open

DO NOT exceed 3 m between bait stations for mice, or 9 m for rats in Clean-Out treatments

DO check for pellets pulled by rodents into the open

### **Baiting Strategy**

- Eliminate as far as practical, all alternative food sources
- Bait infested area. Place bait stations under cover.
- Set out bait stations in places frequented by mice and rats in and around buildings (within 2 m) or enclosed spaces, eg drains
- Actively search for, remove and destroy dead and moribund rodents whilst bait is being used

### ***Clean-Out Treatments (Pulse Baiting)***

- Clean-out treatments are used to remove existing infestations of rats and mice
- Always record the number and locations of bait stations for the purpose of bait replacement and later removal
- For rats, place 1 to 3 bait stations at intervals of 5 to 9 m in infested areas
- For mice, place single bait stations at 2 to 3 m intervals in infested areas
- Place bait stations so as to prevent access to non-target animals. Fix the stations securely to the substrate with nails or wire, where necessary.
- Use the greater number of bait stations and shorter distance between bait placements when infestations are heavy
- Conduct initial inspection of bait points after 3 or 4 days. Replace bait stations where the bait has been consumed. Repeat approximately 4 days later and then weekly. Replace contaminated or spoiled baits.
- Baiting for at least 2 weeks (3 to 4 pulses) will be necessary to reduce rat/mouse numbers to a low level. Although heavy infestations may require longer treatments to achieve complete eradication.
- Discontinue the treatment when effective control has been achieved and remove all bait and bait stations unless a maintenance treatment is to be undertaken

### ***Monitoring and/or Maintenance Treatments***

- Monitoring and maintenance treatments are required at sensitive sites where there is no current rodent infestation but where the prevention of such infestation is essential to prevent significant risks to human health. Such treatments may be a condition for audit approval within food safety guidelines. However, all use of this product must comply with the conditions of this label.
- Always record the number and locations of bait stations for the purpose of bait replacement and later removal
- For rats and mice place one station at intervals of between 15 and 30 m in areas prone to rodent infestation
- Place bait stations so as to prevent access to non-target animals. Always use tamper-resistant bait stations for monitoring and maintenance treatments and fix with nails or wire, if necessary.
- Use placebo or unpoisoned bait blocks if the placement of the bait away from buildings presents a hazard to non-target animals
- When a rodent infestation is detected the operator should revert to the directions for clean-out treatments

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

## **GENERAL INSTRUCTIONS**

Most deaths occur 4 to 7 days after consumption of bait. Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Record the number and location of bait stations for the purpose of replenishment and later removal.

**#Note:** One feed is sufficient to kill rats and mice. Rats and mice will continue to eat the bait after a lethal dose has been eaten.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Depending on the amount eaten, this bait may kill or injure dogs, cats, pigs, poultry and wildlife. **DO NOT** place baits in locations which are accessible to domestic animals, livestock or birds. **DO NOT** contaminate dams, waterways and drains with the bait or its used container. The use of this product for the control of protected native rodents and rodent like animals requires permission from wildlife authorities. The product is restricted to use in and around buildings (within 2 m) or enclosed spaces, eg drains.

## **STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight. Keep out of reach of children and pets. Dispose of empty containers and unused or untaken baits by burying in an approved dump. Look for dead rats or mice, burn or bury them, **DO NOT** place in refuse bins or rubbish tips. Ensure baits are removed when rodent activity has ceased.

## **SAFETY DIRECTIONS**

**Poisonous if swallowed. DO NOT touch bait. If on skin and after each baiting, wash thoroughly with soap and water. Wash hands after use.**

## **FIRST AID**

**If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126. Vitamin K1 (Phytomenadione) is antidotal.**

## **NOTE TO PHYSICIANS AND VETERINARIANS**

**Brodifacoum, the active constituent of TALON Pellets, is a long lasting anticoagulant chemical, which if ingested by humans, domestic animals or pets, can reduce the clotting power of the blood and haemorrhage may result. Vitamin K1 (Phytomenadione) should be administered urgently by medical, appropriate para-medical personnel or veterinarians. If administered intravenously the injection must be given slowly.**

## **SAFETY DATA SHEET**

If additional hazard information is required refer to the Safety Data Sheet. For a copy phone 1800 067 108, or visit our website at [www.syngenta.com.au](http://www.syngenta.com.au)

## **DISCLAIMER**

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.

Product names marked ® or ™, the ALLIANCE FRAME  
the SYNGENTA Logo and the PURPOSE ICON  
are Trademarks of a Syngenta Group Company

