

Product Name: DITRAC RODENTICIDE
 APVMA Approval No: 49868/113073



Label Name:	DITRAC RODENTICIDE
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	0.05 g/kg BRODIFACOUM contains Denatonium Benzoate (Bittering Agent)
Mode of Action:	
Statement of Claims:	Easy to use One Feed - Fast Acting
Net Contents:	NET 2.5 kg PRIMARY PACK CONTAINS 100 X 25 g BAIT STATIONS WHICH ARE ILLEGAL TO SELL SEPARATELY DO NOT DESTROY OUTER CONTAINER WHILE PRODUCT STILL REMAINS NET 5 kg PRIMARY PACK CONTAINS 100 X 50 g BAIT STATIONS WHICH ARE ILLEGAL TO SELL SEPARATELY DO NOT DESTROY OUTER CONTAINER WHILE PRODUCT STILL REMAINS NET 25 g, 50 g, 10 kg
Restrains:	DO NOT exceed 3 m between bait stations for mice, or 9 m for rats. DO NOT use bait stations further than 2 m from buildings.
Directions for Use:	DIRECTIONS FOR USE [2.5 kg(100 x 25 g), 5 kg (100 x 50 g), 10 kg Mice require several bait stations, 50 g per bait station (less than 1 g will kill a mouse). Rats require large quantities of bait in one or two locations, 50 - 100 g per station (4 - 6 g will kill a rat). For effective control continue until feeding ceases.

	<p>Baiting strategy: Eliminate as far as practicable, all other food sources. Bait areas where conditions might encourage rodent activity. Place bait stations under cover. Inspect baits regularly[during routine inspections]. Replace eaten or old/deteriorated baits. If eaten quickly increase number of bait stations. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.</p> <p>DIRECTIONS FOR USE 25 g and 50 g bait stations The DITRAC 25 g [or 50 g] pack is a ready to use 'bait station'. The feeding behaviour of mice is different to that of rats. Rats require a larger quantity of bait. Place bait stations in areas of known rodent activity.</p> <p>Baiting Strategy: Eliminate as far as practicable, all other food sources. Bait areas where conditions might encourage rodent activity. Place baits under cover. Inspect baits regularly[during routine inspections]. Replace eaten or old/deteriorated baits. If eaten quickly increase number of bait stations. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.</p> <p>NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.</p>
--	---

Other Limitations:	THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.
--------------------	--

Withholding Periods:	
----------------------	--

Trade Advice:	
---------------	--

General Instructions:	<p>GENERAL INSTRUCTIONS Pellets are not subject to wind dispersal. Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail or fix in place if relevant. Record the number and locations of bait stations for the purpose of replenishment and later removal. Most deaths occur 4 to 7 days after consumption of bait.</p>
-----------------------	--

Resistance Warning:	
---------------------	--

Precautions:	<p>KEEP OUT OF REACH OF CHILDREN. Keep away from food and water and surfaces which come in contact with food and water.</p>
--------------	---

Protections:	<p>PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate dams, streams, rivers, watercourses or drains with the bait or used containers. The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife and care should be exercised in its use. DO NOT place baits in locations which are accessible to domestic animals, livestock or birds. The product is restricted to use in and around buildings, (within 2 metres) or enclosed spaces.</p>
--------------	--

Storage and Disposal:	<p>Store in the tightly closed, original container in a dry well-ventilated area, as cool as possible out of direct sunlight. Shake empty container onto baiting site. Break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of</p>
-----------------------	---

waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product. Search for and burn or bury all rodent bodies, do not place in refuse bins or in rubbish tips. Ensure uneaten baits are removed and disposed of with the empty containers when the rodent activity has ceased.

Safety Directions:

Poisonous if swallowed. Do not inhale dust. Do not touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water. Vitamin K1 [phytomenadione] is antidotal.

First Aid Instructions:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26; New Zealand 0800 764 766.

First Aid Warnings: